

Online Supplement

Screening Methods

Based on at least two-view screen-film (n=4351) or digital (n=3122) mammography, visually estimated breast density was recorded as < 25%; 26 to 40%; 41 to 60%; 61 to 80%; or > 80% dense. All radiologist investigators met experience requirements and had completed qualification tasks as previously detailed.⁷ Computer-assisted detection was not permitted. Ultrasound scanning was physician performed within two weeks of the mammogram (and usually the same day), with standardized technique and documentation,⁷ using high-resolution linear array broad bandwidth transducers with maximum frequency of at least 12 MHz. Interpretive criteria for both mammography and ultrasound were standardized as well.³⁵ For 1.6% of analyzable participants, the first study screen was their first-ever (baseline) mammographic examination;⁷ otherwise, prior breast imaging was available at the time of interpretation for all screens.

MR Technique

To minimize background parenchymal enhancement, MR examinations were scheduled within 7 to 14 days of onset of menses in premenopausal women when possible and prior to any biopsies prompted by screening mammography or ultrasound. Simultaneous bilateral contrast-enhanced breast MRI utilized dedicated breast coils on 1.5T systems. Axial or

sagittal T1 and fat-suppressed T2 or inversion recovery images were obtained prior to contrast injection. A three-dimensional spoiled gradient echo volume acquisition with fat suppression was obtained through both breasts both prior to and a minimum of three times following intravenous power injection of 0.1 mmol/kg Gd-DTPA, with entirety of both breasts imaged within 3 minutes, delayed imaging for minimum total of 6 minutes, and at least 3 post-contrast acquisitions through both breasts. In-plane voxel size was not greater than 1.0-mm x 1.0-mm x 3-mm slice thickness. Images were viewed with subtraction technique and maximum intensity projection technique. Prior to accruing MRI participants, sites submitted a deidentified sample contrast-enhanced MRI examination for review by Dr. R. Edward Hendrick for quality assurance and to adjust scanning parameters. Computer-assisted kinetic analysis could be performed.

Features and assessments of each lesion and by breast were recorded using BI-RADS.^{14,36,37} To allow meaningful receiver operating characteristic (ROC) curve analysis, we did not allow use of BI-RADS 0, incomplete.

Recommendations for routine annual imaging, short interval follow-up in 6 months, additional imaging, and/or biopsy, were recorded separately.

Additional Results

Time between Screens and Time to Perform US

Time between screening mammography rounds averaged 378 to 380 days (Table E1) and between 24-month mammogram and protocol MRI, 22 days. Time to perform screening ultrasound decreased from 19 minutes at time 0 to less than 15 minutes at time 24 months (Table E2).

Supplemental Yield of US after Digital Mammography

Across three years of screening, supplemental US provided significant detection benefit with both film and digital mammography, with supplemental yield of 3.7 per 1000 (95%CI 2.1 to 5.5) for film and 5.1 per 1000 (95%CI 2.7 to 7.7) for digital mammography ($p<.001$ each), and for all women less than age 50 (3.8 per 1000, 95%CI 1.4 to 6.5) or at least 50 years old (4.5 per 1000, 95%CI 2.8 to 6.1) ($p=.003$ and $p<.001$ vs. mammography alone, respectively).

References cited in online supplement

1. Buchberger W, Niehoff A, Obrist P, DeKoekkoek-Doll P, Dunser M. Clinically and mammographically occult breast lesions: detection and classification with high-resolution sonography. *Semin Ultrasound CT MR*. 2000;21(4):325-336.
2. Crystal P, Strano SD, Shcharynski S, Koretz MJ. Using sonography to screen women with mammographically dense breasts. *AJR Am J Roentgenol*. Jul 2003;181(1):177-182.
3. Gordon PB, Goldenberg SL. Malignant breast masses detected only by ultrasound. A retrospective review [see comments]. *Cancer*. 1995;76(4):626-630.

4. Kaplan SS. Clinical utility of bilateral whole-breast US in the evaluation of women with dense breast tissue. *Radiology*. Dec 2001;221(3):641-649.
5. Kolb TM, Lichy J, Newhouse JH. Comparison of the performance of screening mammography, physical examination, and breast US and evaluation of factors that influence them: An analysis of 27,825 patient evaluations. *Radiology*. Oct 2002;225(1):165-175.
6. Leconte I, Feger C, Galant C, et al. Mammography and subsequent whole-breast sonography of nonpalpable breast cancers: the importance of radiologic breast density. *AJR Am J Roentgenol*. Jun 2003;180(6):1675-1679.
7. Berg WA, Blume JD, Cormack JB, et al. Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer. *JAMA*. May 14 2008;299(18):2151-2163.
8. Corsetti V, Ferrari A, Ghirardi M, et al. Role of ultrasonography in detecting mammographically occult breast carcinoma in women with dense breasts. *Radiol Med (Torino)*. Apr 2006;111(3):440-448.
9. Tohno E, Ueno E, Watanabe H. Ultrasound screening of breast cancer. *Breast Cancer*. 2009;16(1):18-22.
10. Kuhl C, Weigel S, Schrading S, et al. Prospective multicenter cohort study to refine management recommendations for women at elevated familial risk of breast cancer: the EVA trial. *J Clin Oncol*. Mar 20 2010;28(9):1450-1457.
11. Kuhl CK, Schrading S, Leutner CC, et al. Mammography, breast ultrasound, and magnetic resonance imaging for surveillance of women at high familial risk for breast cancer. *J Clin Oncol*. Nov 20 2005;23(33):8469-8476.
12. Lehman CD, Isaacs C, Schnall MD, et al. Cancer yield of mammography, MR, and US in high-risk women: prospective multi-institution breast cancer screening study. *Radiology*. Aug 2007;244(2):381-388.
13. Sardanelli F, Podo F, D'Agnolo G, et al. Multicenter comparative multimodality surveillance of women at genetic-familial high risk for breast cancer (HIBCRIT study): interim results. *Radiology*. Mar 2007;242(3):698-715.
14. D'Orsi CJ, Bassett LW, Berg WA, et al. *Breast Imaging Reporting and Data System, BI-RADS: Mammography, 4th edition*. Reston: American College of Radiology; 2003.
15. Berg WA, Blume JD, Adams AM, et al. Reasons women at elevated risk of breast cancer refuse breast MR imaging screening: ACRIN 6666. *Radiology*. Jan 2010;254(1):79-87.
16. Leemis LM, Trivedi KS. A comparison of approximate interval estimators for the bernoulli parameter. *Amer Statistician*. 1996;50:63-68.
17. Fleiss JL, Levin B, Paik MC. *Statistical Methods for Rates and Proportions*. Hoboken: Wiley-Interscience; 2003.
18. Efron B. Bootstrap Methods: Another Look at the Jackknife. *Ann Stat*. 1979;7(1):1-26.
19. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*. Sep 1988;44(3):837-845.
20. Nelson HD, Tyne K, Naik A, Bougatsos C, Chan BK, Humphrey L. Screening for breast cancer: an update for the U.S. Preventive Services Task Force. *Ann Intern Med*. Nov 17 2009;151(10):727-737, W237-742.
21. Schell MJ, Yankaskas BC, Ballard-Barbash R, et al. Evidence-based target recall rates for screening mammography. *Radiology*. Jun 2007;243(3):681-689.

22. Kriege M, Brekelmans CT, Boetes C, et al. Differences between first and subsequent rounds of the MRISC breast cancer screening program for women with a familial or genetic predisposition. *Cancer*. Jun 1 2006;106(11):2318-2326.
23. Leach MO, Boggis CR, Dixon AK, et al. Screening with magnetic resonance imaging and mammography of a UK population at high familial risk of breast cancer: a prospective multicentre cohort study (MARIBS). *Lancet*. May 21-27 2005;365(9473):1769-1778.
24. Warner E, Plewes DB, Hill KA, et al. Surveillance of BRCA1 and BRCA2 mutation carriers with magnetic resonance imaging, ultrasound, mammography, and clinical breast examination. *Jama*. Sep 15 2004;292(11):1317-1325.
25. Schleinitz MD, De Palo DM, Blume J, D., Cormack JB, Pisano ED, Berg WA. Tolerability of breast cancer screening, diagnostic, and biopsy procedures: Implications for new screening modalities. An ACRIN 6666 Substudy. *Quality of Life Res*. 2011;submitted.
26. Saslow D, Boetes C, Burke W, et al. American cancer society guidelines for breast screening with MRI as an adjunct to mammography. *CA Cancer J Clin*. Mar-Apr 2007;57(2):75-89.
27. Berg WA. Tailored supplemental screening for breast cancer: what now and what next? *AJR Am J Roentgenol*. Feb 2009;192(2):390-399.
28. Brennan ME, Houssami N, Lord S, et al. Magnetic resonance imaging screening of the contralateral breast in women with newly diagnosed breast cancer: systematic review and meta-analysis of incremental cancer detection and impact on surgical management. *J Clin Oncol*. Nov 20 2009;27(33):5640-5649.
29. Brennan S, Liberman L, Dershaw DD, Morris E. Breast MRI screening of women with a personal history of breast cancer. *AJR Am J Roentgenol*. Aug 2010;195(2):510-516.
30. Demartini WB, Kalish GM, Peacock S, Eby PR, Gutierrez RL, Lehman CD. Screening MRI for high risk women: should patients with a treated personal history of breast cancer be screened? Paper presented at: Radiologic Society of North America; November 28, 2010, 2010; Chicago, IL.
31. http://www.cms.gov/PFSlookup/01_Overview.asp#TopOfPage accessed September 8, 2010.
32. Smith RA, Duffy SW, Gabe R, Tabar L, Yen AM, Chen TH. The randomized trials of breast cancer screening: what have we learned? *Radiol Clin North Am*. Sep 2004;42(5):793-806, v.
33. Ohuchi N, Suzuki A, Sakurai Y, et al. Current Status and Problems of Breast Cancer Screening. *JMAJ*. 2009;52:45-49.
34. Sardanelli F, Podo F, Santoro F, et al. Multicenter surveillance of women at high genetic breast cancer risk using mammography, ultrasonography, and contrast-enhanced magnetic resonance imaging (the high breast cancer risk Italian 1 study): final results. *Invest Radiol*. Feb 2011;46(2):94-105.
35. <http://acrin.org/Portals/0/Protocols/6666/Protocol-ACRIN%206666%20Admin%20Update%2011.30.07.pdf>.
36. Mendelson EB, Baum JK, Berg WA, Merritt CRB, Rubin E. *Breast Imaging Reporting and Data System, BI-RADS: Ultrasound, 1st Ed*. Reston: American College of Radiology; 2003.
37. Ikeda DM, Hylton NM, Kuhl CK, et al. *Breast Imaging Reporting and Data System, BI-RADS: Magnetic Resonance Imaging*. Reston: American College of Radiology; 2003.

CONFIDENTIAL IN PRESS TO JAMA

Table E1. Time between Screens (days)

	Year 1 mammo to year 2 mammo	Year 2 mammo to year 3 mammo	Year 3 mammo to protocol MRI
N	2490	2284	612
mean	380	378	25
SD	32	33.9	18.7
median	371	371	22
range	269-675	231-683	0-87

Table E2. Time to Perform Screening US (minutes)

	Year 1	Year 2	Year 3
N	2659	2493 ^a	2321
mean	19.2	16.7	14.7
SD	11.9	10.4	9.2
median	17	15	13

	Year 1	Year 2	Year 3
range	1-144	1-166	1-121

^a Three participants not included in year 1 analysis set were included in year 2.

1 **Table 5A. Results of Supplemental US Screening in Addition to Mammography for Women with and without Personal History of Breast**
2 **Cancer (PHBC)**

3

		Mammography (M) alone			Combined Mammography and US (M + US)			Difference of (M + US) and M alone			Ultrasound (US) alone		
		Est	(No/ Total)	95% CI	Est	(No/ Total)	95% CI	Est	95% CI	pvalue	Est	(No/ Total)	95% CI
Yield, per 1000 (N/Total)	PHBC	8.2	(33/ 4010)	(5.7, 11.5)	12.5	(50/ 4010)	(9.3, 16.4)	4.2	(2.5, 6.2)	<0.001	8.7	(35/ 4010)	(6.1, 12.1)
	No PHBC	7.5	(26/ 3463)	(4.9, 11.0)	11.8	(41/ 3463)	(8.5, 16.0)	4.3	(2.3, 6.6)	<0.001	6.6	(23/ 3463)	(4.2, 9.9)
	Difference	0.7		(-3.3, 4.4)	0.6		(-4.4, 5.5)	-0.1	(-3.0, 2.9)	0.950	2.1		(-1.9, 5.9)
Sensitivity, % (N/Total)	PHBC	55.9	(33/ 59)	(42.4, 68.8)	84.7	(50/ 59)	(73.0, 92.8)	28.8	(18.6, 40.7)	<0.001	59.3	(35/ 59)	(45.7, 71.9)
	No PHBC	50.0	(26/ 52)	(35.8, 64.2)	78.8	(41/ 52)	(65.3, 88.9)	28.8	(17.3, 40.4)	<0.001	44.2	(23/ 52)	(30.5, 58.7)
	Difference	5.9		(-12.5, 24.6)	5.9		(-8.8, 20.0)	-0.0	(-15.9, 15.6)	0.997	15.1		(-2.4, 32.6)
Specificity, % (N/Total)	PHBC	91.4	(3611/ 3951)	(90.5, 92.3)	83.1	(3283/ 3951)	(81.9, 84.2)	-8.3	(-9.3, -7.4)	<0.001	88.2	(3485/ 3951)	(87.2, 89.2)
	No PHBC	89.4	(3051/ 3411)	(88.4, 90.5)	77.8	(2654/ 3411)	(76.4, 79.2)	-11.6	(-12.8, -10.4)	<0.001	84.0	(2865/ 3411)	(82.7, 85.2)
	Difference	1.9		(0.7, 3.3)	5.3		(3.5, 7.1)	3.3	(1.7, 4.9)	<0.001	4.2		(2.7, 5.7)
Recall Rate, % (N/Total)	PHBC	9.3	(373/ 4010)	(8.4, 10.2)	17.9	(718/ 4010)	(16.7, 19.1)	8.6	(7.7, 9.5)	<0.001	12.5	(501/ 4010)	(11.5, 13.6)
	No PHBC	11.1	(386/ 3463)	(10.1, 12.2)	23.0	(798/ 3463)	(21.6, 24.5)	11.9	(10.6, 13.1)	<0.001	16.4	(569/ 3463)	(15.2, 17.7)
	Difference	-1.8		(-3.2, -0.4)	-5.1		(-7.0, -3.4)	-3.3	(-4.9, -1.7)	<0.001	-3.9		(-5.6, -2.4)
PPV1, ^b % (N/Total)	PHBC	8.8	(33/ 373)	(6.2, 12.2)	7.0	(50/ 718)	(5.2, 9.1)	-1.9	(-3.5, -0.2)	0.026	7.0	(35/ 501)	(4.9, 9.6)
	No PHBC	6.7	(26/ 386)	(4.4, 9.7)	5.1	(41/ 798)	(3.7, 6.9)	-1.6	(-3.2, -0.0)	0.050	4.0	(23/ 569)	(2.6, 6.0)
	Difference	2.1		(-1.6, 5.8)	1.8		(-0.3, 4.2)	-0.3	(-2.6, 2.0)	0.808	2.9		(0.8, 6.6)
Short-term	PHBC	2.2	(90/ 4010)	(1.8, 2.8)	7.5	(300/ 4010)	(6.7, 8.3)	5.2	(4.5, 5.9)	<0.001	5.5	(220/ 4010)	(4.8, 6.2)
Follow-up Rate, %	No PHBC	2.0	(70/ 3463)	(1.6, 2.5)	9.4	(324/ 3463)	(8.4, 10.4)	7.3	(6.4, 8.2)	<0.001	7.7	(266/ 3463)	(6.8, 8.6)
(N/Total)	Difference	0.2		(-0.5, 0.9)	-1.9		(-3.1, -0.6)	-2.1	(-3.2, -1.0)	<0.001	-2.2		(-3.3, -1.0)

Biopsy Rate, % (N/Total)	PHBC	2.2	(87/ 4010)	(1.7, 2.7)	6.9	(275/ 4010)	(6.1, 7.7)	4.7	(4.0, 5.3)	<0.001	5.4	(215/ 4010)	(4.7, 6.1)
	No PHBC	2.2	(75/ 3463)	(1.7, 2.7)	9.7	(336/ 3463)	(8.7, 10.7)	7.5	(6.6, 8.4)	<0.001	8.2	(284/ 3463)	(7.3, 9.2)
	Difference	0.0		(-0.6, 0.7)	-2.8		(-4.2, -1.6)	-2.8	(-4.0, -1.8)	<0.001	-2.8		(-4.0, -1.7)
PPV3,^c % (N/Total)	PHBC	36.8	(32/ 87)	(26.7, 47.8)	17.8	(49/ 275)	(13.5, 22.9)	-19.0	(-26.7, -11.7)	<0.001	15.3	(33/ 215)	(10.8, 20.9)
	No PHBC	32.0	(24/ 75)	(21.7, 43.8)	11.0	(37/ 336)	(7.9, 14.9)	-21.0	(-29.1, -13.1)	<0.001	6.7	(19/ 284)	(4.1, 10.3)
	Difference	4.8		(-8.8, 18.6)	6.8		(1.9, 12.2)	2.0	(-8.7, 12.8)	0.714	8.7		(3.7, 14.0)

4

5

6 ^a Yield is the cancer detection rate

7 ^b PPV1 is defined as the malignancy rate among women with a positive screening test (i.e. recalled from screening for further testing or short-
8 interval follow-up).

9 ^c PPV3 is defined as the malignancy rate among women with a positive screening test who underwent biopsy of the same lesion.

10

11 **Table 5B. Results of Supplemental MRI Screening in Addition to Mammography for Women with and without Personal History of Breast**

	M	M+MRI	(M+MRI)-(M)	MRI
--	---	-------	-------------	-----

12 **Cancer (PHBC)**

13

		Est	N/ Total	95% CI	Est	N/ Total	95% CI	Est	95% CI	P	Est	N/ Total	95% CI
Yield ^a per 1000	PHBC	7.3	2/275	(0.9, 26.0)	14.5	4/275	(4.0, 36.8)	7.3	(-6.4, 21.0)	0.500	10.9	3/275	(2.3, 31.5)
	No PHBC	8.9	3/337	(1.8, 25.8)	35.6	12/337	(18.5, 61.4)	26.7	(6.5, 46.9)	0.004	32.6	11/337	(16.4, 57.7)
	Difference	-1.6		(-15.7, 12.4)	-21.1		(-45.7, 3.5)	-19.4	(-40.0, 1.1)	0.063	-21.7		(-44.4, 0.9)
Sensitivity (%)	PHBC	50.0	2/4	(6.8, 93.2)	100.0	4/4	(39.8, 100.0)	50.0	(-24.0, 124.0)	0.500	75.0	3/4	(19.4, 99.4)
	No PHBC	25.0	3/12	(5.5, 57.2)	100.0	12/12	(73.5, 100.0)	75.0	(42.2, 107.8)	0.004	91.7	11/12	(61.5, 99.8)
	Difference	25.0		(-28.4, 78.4)	0.0		(0.0, 0.0)	-25.0	(-78.4, 28.4)	0.359	-16.7		(-63.3, 29.9)
Specificity (%)	PHBC	95.2	258/271	(91.9, 97.4)	78.6	213/271	(73.2, 83.3)	-16.6	(-21.4, -11.8)	<.001	81.5	221/271	(76.4, 86.0)
	No PHBC	89.5	291/325	(85.7, 92.6)	64.0	208/325	(58.5, 69.2)	-25.5	(-30.6, -20.5)	<.001	70.8	230/325	(65.5, 75.7)
	Difference	5.7		(1.5, 9.8)	14.6		(7.4, 21.8)	8.9	(2.4, 15.5)	0.008	10.8		(3.9, 17.6)
PPV1 ^b (%)	PHBC	13.3	2/15	(1.7, 40.5)	6.5	4/62	(1.8, 15.7)	-6.9	(-22.0, 4.8)	0.337	5.7	3/53	(1.2, 15.7)
	No PHBC	8.1	3/37	(1.7, 21.9)	9.3	12/129	(4.9, 15.7)	1.2	(-6.1, 8.3)	0.750	10.4	11/106	(5.3, 17.8)
	Difference	5.2		(-14.3, 24.8)	-2.9		(-10.7, 5.0)	-8.1	(-24.2, 8.1)	0.327	-4.7		(-13.1, 3.7)
Recall (%)	PHBC	5.5	15/275	(3.1, 8.8)	22.5	62/275	(17.7, 27.9)	17.1	(12.3, 21.9)	<.001	19.3	53/275	(14.8, 24.4)
	No PHBC	11.0	37/337	(7.8, 14.8)	38.3	129/337	(33.1, 43.7)	27.3	(22.2, 32.4)	<.001	31.5	106/337	(26.5, 36.7)
	Difference	-5.5		(-9.7, -1.4)	-15.7		(-22.9, -8.5)	-10.2	(-16.8, -3.7)	0.002	-12.2		(-19.2, -5.2)
Short-term Follow-up Rate (%)	PHBC	0.4	1/275	(0.0, 2.0)	14.2	39/275	(10.3, 18.9)	13.8	(9.4, 18.3)	<.001	13.8	38/275	(10.0, 18.5)
	No PHBC	0.6	2/337	(0.1, 2.1)	18.1	61/337	(14.1, 22.6)	17.5	(13.2, 21.9)	<.001	17.5	59/337	(13.6, 22.0)
	Difference	-0.2		(-1.3, 0.9)	-3.9		(-9.9, 2.0)	-3.7	(-9.6, 2.2)	0.219	-3.7		(-9.6, 2.2)
Biopsy Rate (%)	PHBC	1.5	4/275	(0.4, 3.7)	5.5	15/275	(3.1, 8.8)	4.0	(1.3, 6.7)	<.001	4.0	11/275	(2.0, 7.0)
	No PHBC	1.8	6/337	(0.7, 3.8)	13.1	44/337	(9.6, 17.1)	11.3	(7.6, 14.9)	<.001	12.2	41/337	(8.9, 16.1)

	Difference	-0.3		(-2.3, 1.6)	-7.6		(-12.2, -3.0)	-7.3	(-11.4, -3.1)	<.001	-8.2		(-12.4, -3.9)
PPV3^c (%)	PHBC	50.0	2/4	(6.8, 93.2)	26.7	4/15	(7.8, 55.1)	-23.3	(-66.7, 20.0)	0.251	18.2	2/11	(2.3, 51.8)
	No PHBC	50.0	3/6	(11.8, 88.2)	25.0	11/44	(13.2, 40.3)	-25.0	(-58.3, 10.6)	0.166	24.4	10/41	(12.4, 40.3)
	Difference	0.0		(-63.4, 63.4)	1.7		(-22.5, 25.8)	1.7	(-51.9, 55.2)	0.951	-6.2		(-31.5, 19.1)

14

15

16

17

18 ^a Yield is the cancer detection rate

19 ^b PPV1 is defined as the malignancy rate among women with a positive screening test (i.e. recalled from screening for further testing or short-
20 interval follow-up).

21 ^c PPV3 is defined as the malignancy rate among women with a positive screening test who underwent biopsy of the same lesion.

22

23

24

25